Large-Type Edition



The University of the State of New York REGENTS HIGH SCHOOL EXAMINATION

ALGEBRAII

Thursday, January 25, 2024 — 1:15 to 4:15 p.m., only

Student Name _____

School Name _____

The possession or use of any communications device is strictly prohibited when taking this examination. If you have or use any communications device, no matter how briefly, your examination will be invalidated and no score will be calculated for you.

Print your name and the name of your school on the lines above.

A separate answer sheet for **Part I** has been provided to you. Follow the instructions from the proctor for completing the student information on your answer sheet.

This examination has four parts, with a total of 37 questions. You must answer all questions in this examination. Record your answers to the Part I multiple-choice questions on the separate answer sheet. Write your answers to the questions in **Parts II**, **III**, and **IV** directly in this booklet. All work should be written in pen, except graphs and drawings, which should be done in pencil. Clearly indicate the necessary steps, including appropriate formula substitutions, diagrams, graphs, charts, etc. Utilize the information provided for each question to determine your answer. Note that diagrams are not necessarily drawn to scale.

Algebra II – Jan. '24

The formulas that you may need to answer some questions in this examination are found at the end of the examination. You may remove this sheet from this booklet.

Scrap paper is not permitted for any part of this examination, but you may use the blank spaces in this booklet as scrap paper. A sheet of scrap graph paper is provided at the end of this booklet for any question for which graphing may be helpful but is not required. You may remove this sheet from this booklet. Any work done on this sheet of scrap graph paper will not be scored.

When you have completed the examination, you must sign the statement printed at the end of the answer sheet, indicating that you had no unlawful knowledge of the questions or answers prior to the examination and that you have neither given nor received assistance in answering any of the questions during the examination. Your answer sheet cannot be accepted if you fail to sign this declaration.

Notice ...

A graphing calculator and a straightedge (ruler) must be available for you to use while taking this examination.

DO NOT OPEN THIS EXAMINATION BOOKLET UNTIL THE SIGNAL IS GIVEN.

Part I

Answer all 24 questions in this part. Each correct answer will receive 2 credits. No partial credit will be allowed. Utilize the information provided for each question to determine your answer. Note that diagrams are not necessarily drawn to scale. For each statement or question, choose the word or expression that, of those given, best completes the statement or answers the question. Record your answers on your separate answer sheet. [48]

1 A cafeteria food manager studied the lunchtime eating habits of a group of employees in their office building. The purpose of the study was to determine the proportion of employees who purchased lunch in the cafeteria, brought their lunch from home, or purchased lunch from an outside vendor. This collection of data would best be classified as

Use this space for computations.

(1) a census


(3) an observational study

(2) an experiment

(4) a simulation



Use this space for computations.



Algebra II – Jan. '24


3 Given 3 is a root of $f(x) = x^4 - x^3 - 21x^2 + 45x$, what are the other unique roots of f(x)?

Use this space for computations.

- (1) -5, only (3) -3, 1 and 5
- (2) -5 and 0 (4) -5, -3 and 0

4 Given $p \neq q, p = \left(\frac{1}{2}\right)^q$, expressed in logarithmic form, is equivalent to (1) $\log_p\left(\frac{1}{2}\right) = q$ (3) $\log_1(p) = q$ (2) $\log_q(p) = \frac{1}{2}$ (4) $\log_1(q) = p$ **5** Which graph best represents the graph of $f(x) = (x + a)^2(x - b)$, where *a* and *b* are positive real numbers?

Use this space for computations.



Algebra II - Jan. '24

[OVER]

6 The equations y = 3t + 6 and $y = (1.82)^t$ approximately model the growth of two separate populations where t > 0. What is the best approximation of the time, t, at which the populations are the same?

Use this space for computations.

- (1) -1.9 (3) 5.1
- (2) 0.3 (4) 21.3

- **7** Given y = -2x and $x^2 + y^2 = 5$, the point of intersection in Quadrant II is
 - (1) (1, -2) (3) (-1, 1)
 - (2) (-2, 1) (4) (-1, 2)

8 The rational expression $\frac{2x^4 - 5x^2 + 3x - 2}{x - 3}$ is equivalent to (1) $2x^3 - 5x - 12 - \frac{38}{x - 3}$ (2) $2x^3 + 6x^2 + 13x + 42 + \frac{124}{x - 3}$ (3) $2x^3 - 5x + 18 - \frac{56}{x - 3}$ (4) $2x^3 - 6x^2 + 13x - 36 + \frac{106}{x - 3}$

9 The equation of the parabola that has its focus at the point (-3, 2) and directrix at y = 0 is

(1)
$$y = \frac{1}{4}(x+3)^2 + 1$$

(3) $y = \frac{1}{8}(x+3)^2 + 1$
(2) $y = \frac{1}{4}(x-3)^2 + 1$
(4) $y = \frac{1}{8}(x-3)^2 + 1$

Algebra II – Jan. '24

Use this space for

computations.

 10 The seventh term of the geometric sequence $\sqrt{6}, -2\sqrt{3}, 2\sqrt{6}, -4\sqrt{3}...$ is

 (1) $6\sqrt{6}$ (3) $8\sqrt{6}$

 (2) $-6\sqrt{3}$ (4) $-8\sqrt{3}$

Use this space for computations.

11 A company wishes to determine the cooking time for one pound of spaghetti. The company's technicians cooked one pound of spaghetti and recorded the time needed for the spaghetti to be ready to eat. Repeating this process 35 times resulted in an approximately normal distribution, with a mean of 9.82 minutes and a standard deviation of 1.4 minutes. In which interval should the middle 95% of cooking times fall?

(1) (8.42, 11.22)	(3) (9.35, 10.29)
(2) $(7.02, 12.62)$	(4) (6.82, 11.32)

12 Given $f(x) = 2x^2 + 7x - 15$ and g(x) = 3 - 2x, what is $\frac{f(x)}{g(x)}$ for all defined values? (1) -x - 5 (3) x - 5 Use this space for computations.

(1)
$$-x - 5$$
 (3) $x - 5$
(2) $-x + 5$ (4) $x + 5$

13 Which equation is equivalent to
$$P = 210x^{\frac{4}{3}}y^{\frac{7}{3}}$$
?

(1)
$$P = \sqrt[3]{210x^4y^7}$$
 (3) $P = 210xy^2\sqrt[3]{xy}$
(2) $P = 70xy^2\sqrt[3]{xy}$ (4) $P = 210xy^2\sqrt[3]{x^3y^5}$

14 The average cost of a gallon of milk in the United States between the years of 1995 and 2018 can be modeled by the equation $P(t) = -0.0004t^3 + 0.0114t^2 - 0.0150t + 2.6602$, where P(t)represents the cost, in dollars, and t is time in years since January 1995. During this time period, in what year did P(t) reach its maximum?

- (1) 1995 (3) 2014
- $(2) \ 2013 \tag{4} \ 2018$

15 The temperature, F, in degrees Fahrenheit, after t hours of a roast put into an oven is given by the equation $F = 325 - 185e^{-0.4t}$. What was the temperature of the roast when it was put into the oven?

(1) 325	(3) 185
---------	---------

(2) 200 (4) 140

16 The roots of the equation $0 = x^2 + 6x + 10$ in simplest a + bi form are

Use this space for computations.

(1)
$$-3 \pm 2i$$
 (3) $-3 \pm i$

(2) $-6 \pm i$ (4) $-3 \pm i\sqrt{2}$


17 Which equation does *not* represent an identity?

(1)
$$x^2 - y^2 = (x + y)(x - y)$$

(2) $(x - y)^2 = (x - y)(x - y)$
(3) $(x + y)^2 = x^2 + 2xy + y^2$
(4) $(x + y)^3 = x^3 + 3xy + y^3$

18 Two surveys were conducted to estimate the proportion of teens who use social media at least once per day.

Use this space for computations.



Question 18 is continued on the next page.

Question 18 continued

Use this space for computations.

Based on these results, it was determined that approximately 75% of teens use social media at least once per day. What is the best explanation of the difference in the results between the two surveys?

- (1) The smaller sample size of five teens resulted in a smaller margin of error and should provide a more accurate estimate.
- (2) The smaller sample size of five teens resulted in a bigger margin of error and should provide a more accurate estimate.
- (3) The larger sample size of 50 teens resulted in a smaller margin of error and should provide a more accurate estimate.
- (4) The larger sample size of 50 teens resulted in a bigger margin of error and should provide a more accurate estimate.

19 Given $f(x) = x^3 - 3$ and $f^{-1}(x) = \sqrt[3]{x - 3b}$, the value of *b* is (1) 1
(3) 3
(2) -1
(4) -3

Algebra II – Jan. '24

20 Robert is buying a car that costs \$22,000. After a down payment of \$4000, he borrows the remainder from a bank, a six year loan at 6.24% annual interest rate. The following formula can be used to calculate his monthly loan payment.

$$R = \frac{(P)(i)}{1 - (1 + i)^{-t}}$$

R = monthly payment P = loan amount i = monthly interest ratet = time, in months

Robert's monthly payment will be

(1) \$298.31
(2) \$300.36
(3) \$307.35
(4) \$367.10

Use this space for computations.

21 Given $\tan \theta = -\frac{4}{3}$ where $\frac{\pi}{2} < \theta < \pi$, what is the value of sec θ ? (1) $-\frac{5}{3}$ (3) $\frac{4}{5}$ (2) $-\frac{3}{5}$ (4) $\frac{5}{3}$

22 To solve the equation $\frac{7}{x+7} + \frac{4x}{x-7} = \frac{3x+7}{x-7}$, Joan's first step is to multiply both sides by the least common denominator. Which statement is true?

- (1) -14 is an extraneous solution.
- (2) 7 and -7 are extraneous solutions.
- (3) 7 is an extraneous solution.
- (4) There are no extraneous solutions.

Use this space for

computations.

23 Beginning July 1, 2019, Michelle deposited \$250 into an account that yields 0.15% each month. She continued to make \$250 deposits into this account on the first of each month for 3 years. Which expression represents the amount of money that was in the account after her last deposit was made on June 1, 2022?

(1)
$$250(1.0015)^3$$

(2) $250(1.0015)^{36}$
(3) $\frac{250 - 250(1.0015)^3}{1 - 1.0015}$
(4) $\frac{250 - 250(1.0015)^{36}}{1 - 1.0015}$

24 A study of the red tailed hawk population in a given area shows the population, H(t), can be represented by the function $H(t) = 50(1.19)^t$ where t represents the number of years since the study began. In terms of the monthly rate of growth, the population can be best approximated by the function

(1) $H(t) = 50(1.015)^{12t}$ (3) $H(t) = 50(1.19)^{12t}$ (2) $H(t) = 50(1.15)^{\frac{t}{12}}$ (4) $H(t) = 50(1.19)^{\frac{t}{12}}$

Use this space for computations.

GO RIGHT ON TO THE NEXT PAGE \Longrightarrow

Part II

Answer all 8 questions in this part. Each correct answer will receive 2 credits. Clearly indicate the necessary steps, including appropriate formula substitutions, diagrams, graphs, charts, etc. Utilize the information provided for each question to determine your answer. Note that diagrams are not necessarily drawn to scale. For all questions in this part, a correct numerical answer with no work shown will receive only 1 credit. All answers should be written in pen, except for graphs and drawings, which should be done in pencil. [16]

25 Factor $x^3 + 4x^2 - 9x - 36$, completely.


Work space for question 25 is continued on the next page.

Question 25 continued

26 Determine if x + 4 is a factor of $2x^3 + 10x^2 + 4x - 16$. Explain your answer.

27 An initial investment of \$1000 reaches a value, V(t), according to the model $V(t) = 1000(1.01)^{4t}$, where t is the time in years.

Determine the average rate of change, to the *nearest dollar per year*, of this investment from year 2 to year 7.



29 The heights of the members of a ski club are normally distributed. The average height is 64.7 inches with a standard deviation of 4.3 inches. Determine the percentage of club members, to the *nearest percent*, who are between 67 inches and 72 inches tall.

30 The explicit formula $a_n = 6 + 6n$ represents the number of seats in each row in a movie theater, where *n* represents the row number. Rewrite this formula in recursive form.

31 Express $(2xi^3 - 3y)^2$ in simplest form.

32 A survey was given to 1250 randomly selected high school students at the end of their junior year. The survey offered four post-graduation options: two-year college, four-year college, military, or work. Of the 1250 responses, 475 chose a four-year college. State *one* possible conclusion that can be made about the population of high school juniors, based on this survey.

Work space for question 32 is continued on the next page.

Question 32 continued

Part III

Answer all 4 questions in this part. Each correct answer will receive 4 credits. Clearly indicate the necessary steps, including appropriate formula substitutions, diagrams, graphs, charts, etc. Utilize the information provided for each question to determine your answer. Note that diagrams are not necessarily drawn to scale. For all questions in this part, a correct numerical answer with no work shown will receive only 1 credit. All answers should be written in pen, except for graphs and drawings, which should be done in pencil. [16]

33 A researcher wants to determine if nut allergies and milk allergies are related to each other. The researcher surveyed 1500 people and asked them if they are allergic to nuts or milk. The survey results are summarized in the table below.

	Allergic to Nuts	Not Allergic to Nuts
Allergic to Milk	3	42
Not Allergic to Milk	12	1443

Question 33 is continued on the next page.

Question 33 continued

Determine the probability that a randomly selected survey respondent is allergic to milk.

Determine the probability that a randomly selected survey respondent is allergic to milk, given that the person is allergic to nuts.

Based on the survey data, determine whether nut allergies and milk allergies are independent events. Justify your answer.

34 Algebraically solve for $x: 2x = 6 + 2\sqrt{x-1}$

Work space for question 34 is continued on the next page.


Question 34 continued

35 During the summer, Adam saved \$4000 and Betty saved \$3500. Adam deposited his money in Bank *A* at an annual rate of 2.4% compounded monthly. Betty deposited her money in Bank *B* at an annual rate of 4% compounded quarterly. Write two functions that represent the value of each account after *t* years if no other deposits or withdrawals are made, where Adam's account value is represented by A(t), and Betty's by B(t).

Question 35 continued

Using technology, determine, to the *nearest tenth of a year*, how long it will take for the two accounts to have the same amount of money in them. Justify your answer.

36 On the graph below, draw *at least one* complete cycle of a sine graph passing through point (0,2) that has an amplitude of 3, a period of π , and a midline at y = 2.



Question 36 is continued on the next page.

Algebra II - Jan. '24

Question 36 continued

Based on your graph, state an interval in which the graph is increasing.

Part IV

Answer the question in this part. A correct answer will receive 6 credits. Clearly indicate the necessary steps, including appropriate formula substitutions, diagrams, graphs, charts, etc. Utilize the information provided to determine your answer. Note that diagrams are not necessarily drawn to scale. A correct numerical answer with no work shown will receive only 1 credit. All answers should be written in pen, except for graphs and drawings, which should be done in pencil. [6]

37 A manufacturer of sweatshirts finds that profits and costs fluctuate depending on the number of products created. Creating more products doesn't always increase profits because it requires additional costs, such as building a larger facility or hiring more workers. The manufacturer determines the profit, p(x), in thousands of dollars, as a function of the number of sweatshirts sold, x, in thousands. This function, p, is given below.

 $p(x) = -x^3 + 11x^2 - 7x - 69$

Graph y = p(x), over the interval $0 \le x \le 9$, on the set of axes on the next page.

Question 37 is continued on the next page.

Question 37 continued У -30-≻X 1 Question 37 is continued on the next page.

Algebra II – Jan. '24


Question 37 continued


Over the given interval, state the coordinates of the maximum of p and round all values to the *nearest integer*. Explain what this point represents in terms of the number of sweatshirts sold and profit.

Question 37 is continued on the next page.

Question 37 continued

Determine how many sweatshirts, to the *nearest whole sweatshirt*, the manufacturer would need to produce in order to first make a positive profit. Justify your answer.





High School Math Reference Sheet

1 inch = 2.54 centimeters	1 kilometer = 0.62 mile	1 cup = 8 fluid ounces
1 meter = 39.37 inches	1 pound = 16 ounces	1 pint = 2 cups
1 mile = 5280 feet	1 pound = 0.454 kilogram	1 quart = 2 pints
1 mile = 1760 yards	1 kilogram = 2.2 pounds	1 gallon = 4 quarts
1 mile = 1.609 kilometers	1 ton = 2000 pounds	1 gallon = 3.785 liters
	_	1 liter = 0.264 gallon

1 liter = 1000 cubic centimeters

Triangle	$A = \frac{1}{2}bh$	Pythagorean Theorem	$a^2 + b^2 = c^2$
Parallelogram	A = bh	Quadratic Formula	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Circle	$A = \pi r^2$	Arithmetic Sequence	$a_n = a_1 + (n-1)d$
Circle	$C = \pi d \text{ or } C = 2\pi r$	Geometric Sequence	$a_n = a_1 r^{n-1}$
General Prisms	V = Bh	Geometric Series	$S_n = \frac{a_1 - a_1 r^n}{1 - r}$ where $r \neq 1$

The Reference Sheet is continued on the next page.

Reference Sheet — concluded

Cylinder	$V = \pi r^2 h$
Sphere	$V = \frac{4}{3}\pi r^3$
Cone	$V = \frac{1}{3}\pi r^2 h$
Pyramid	$V = \frac{1}{3}Bh$

Radians	1 radian = $\frac{180}{\pi}$ degrees
Degrees	1 degree = $\frac{\pi}{180}$ radians
Exponential Growth/Decay	$A = A_0 e^{k(t - t_0)} + B_0$